

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 1-[5-(4-Bromophenyl)-3-(4-fluorophenyl)-4,5-dihydro-1H-pyrazol-1-yl]-ethanone

Hoong-Kun Fun,<sup>a,\*</sup> Wan-Sin Loh,<sup>a,§</sup> M. Sapnakumari,<sup>b</sup> B. Narayana<sup>b</sup> and B. K. Sarojini<sup>c</sup>

<sup>a</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, <sup>b</sup>Department of Studies in Chemistry, Mangalore University, Mangalagangotri 574 199, India, and <sup>c</sup>Department of Chemistry, P.A. College of Engineering, Nadupadavu, Mangalore 574 153, India  
Correspondence e-mail: hkfun@usm.my


Received 19 July 2012; accepted 24 July 2012

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}—\text{C}) = 0.002$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.064; data-to-parameter ratio = 26.9.

In the title molecule,  $\text{C}_{17}\text{H}_{14}\text{BrFN}_2\text{O}$ , the benzene rings form dihedral angles of  $6.58$  (6) and  $85.31$  (6)° with the mean plane of the 4,5-dihydro-1H-pyrazole ring (r.m.s. deviation =  $0.0231$  Å). The latter ring is planar with a maximum deviation of  $0.032$  (1) Å. The dihedral angle between the benzene rings is  $78.75$  (6)°. In the crystal, weak  $\text{C}—\text{H} \cdots \text{O}$  and  $\text{C}—\text{H} \cdots \text{F}$  hydrogen bonds link the molecules into corrugated layers parallel to the  $ab$  plane.

## Related literature

For our work on the synthesis of pyrazoline derivatives, see: Samshuddin *et al.* (2011). For related structures, see: Fun *et al.* (2010, 2012). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



## Experimental

## Crystal data

 $\text{C}_{17}\text{H}_{14}\text{BrFN}_2\text{O}$  $M_r = 361.21$ 

Monoclinic,  $P2_1/c$   
 $a = 6.0973$  (5) Å  
 $b = 12.3079$  (11) Å  
 $c = 20.1432$  (16) Å  
 $\beta = 96.700$  (1)°  
 $V = 1501.3$  (2) Å<sup>3</sup>

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 2.75$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.35 \times 0.29 \times 0.12$  mm

## Data collection

Bruker SMART APEXII DUO  
 CCD area-detector  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2009)  
 $T_{\min} = 0.449$ ,  $T_{\max} = 0.735$

20560 measured reflections  
 5389 independent reflections  
 4508 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.064$   
 $S = 1.04$   
 5389 reflections

200 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.50$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.25$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{C4}—\text{H4A} \cdots \text{O1}^{\text{i}}$	0.95	2.45	3.2772 (15)	146
$\text{C14}—\text{H14A} \cdots \text{F1}^{\text{ii}}$	0.95	2.50	3.3806 (15)	153
$\text{C15}—\text{H15A} \cdots \text{O1}^{\text{iii}}$	0.95	2.45	3.3800 (15)	166

Symmetry codes: (i)  $-x, y + \frac{1}{2}, -z + \frac{3}{2}$ ; (ii)  $-x - 1, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iii)  $x - 1, y, z$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and WSL thank Universiti Sains Malaysia (USM) for the Research University Grant (1001/PFIZIK/811160). WSL also thanks the Malaysian Government and USM for the post of Research Officer under the Research University Grant (1001/PFIZIK/811160). BN thanks the UGC for financial assistance through the SAP and BSR one-time grant for the purchase of chemicals.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5325).

## References

- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.  
 Fun, H.-K., Hemamalini, M., Samshuddin, S., Narayana, B. & Yathirajan, H. S. (2010). *Acta Cryst.* **E66**, o582–o583.  
 Fun, H.-K., Quah, C. K., Samshuddin, S., Narayana, B. & Sarojini, B. K. (2012). *Acta Cryst.* **E68**, o975.  
 Samshuddin, S., Narayana, B., Baktir, Z., Akkurt, M. & Yathirajan, H. S. (2011). *Der Pharma Chem.* **3**, 487–493.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

\* Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: C-7581-2009.

## supplementary materials

*Acta Cryst.* (2012). E68, o2586 [doi:10.1107/S1600536812033351]

**1-[5-(4-Bromophenyl)-3-(4-fluorophenyl)-4,5-dihydro-1H-pyrazol-1-yl]ethanone**

**Hoong-Kun Fun, Wan-Sin Loh, M. Sapnakumari, B. Narayana and B. K. Sarojini**

**Comment**

In continuation of our work on the synthesis of pyrazoline derivatives (Fun *et al.*, 2010; Samshuddin *et al.*, 2011), the title compound is prepared and its crystal structure is reported.

In the title compound (Fig. 1), the two benzene rings (C1–C6 & C10–C15) form dihedral angles of 6.58 (6) and 85.31 (6)°, respectively, with the mean plane of 4,5-dihydro-1H-pyrazole ring (N1/N2/C7–C9, r.m.s. deviation = 0.0231 Å). The dihedral angle between the two benzene rings is 78.75 (6)°. Bond lengths and angles are comparable with those in the related structures (Fun *et al.*, 2010, 2012).

In the crystal packing (Fig. 2), intermolecular C—H···O and C—H···F hydrogen bonds (Table 1) link the molecules into corrugated layers parallel to the *ab* plane.

**Experimental**


A mixture of (2*E*)-3-(4-bromophenyl)-1-(4-fluorophenyl)prop-2-en-1-one (3.05 g, 0.01 mol) and hydrazine hydrate (0.48 ml, 0.01 mol) in 30 ml acetic acid was refluxed for 6 h. The reaction mixture was cooled and poured into 50 ml ice-cold water. The precipitate was collected by filtration and purified by recrystallization from ethanol. The single-crystal was grown from acetone by slow evaporation method. *M.p.*: 372–374 K.

**Refinement**

All the H atoms were located geometrically and were refined using a riding model with  $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5 U_{\text{eq}}(\text{C})$  [C–H = 0.95 to 1.00 Å]. A rotating group model was applied to the methyl group. In the final refinement, ten outliers were omitted, namely -2 0 8, 1 4 2, 1 0 4, -2 7 5, -4 2 9, -3 4 9, -1 8 1, -2 2 4, 1 8 0 and 1 1 4, respectively.


**Computing details**

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).



**Figure 1**

The molecular structure of the title compound, showing 50% probability displacement ellipsoids.



**Figure 2**

A portion of the crystal packing viewed along the *a* axis. H atoms not involved in the intermolecular interactions (dashed lines) have been omitted for clarity.

**1-[5-(4-Bromophenyl)-3-(4-fluorophenyl)-4,5-dihydro-1*H*-pyrazol- 1-yl]ethanone**

*Crystal data*

$C_{17}H_{14}BrFN_2O$

$M_r = 361.21$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 6.0973\ (5)\ \text{\AA}$

$b = 12.3079\ (11)\ \text{\AA}$

$c = 20.1432\ (16)\ \text{\AA}$

$\beta = 96.700\ (1)^\circ$

$V = 1501.3\ (2)\ \text{\AA}^3$

$Z = 4$

$F(000) = 728$

$D_x = 1.598\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 7995 reflections

$\theta = 3.3\text{--}32.4^\circ$

$\mu = 2.75\ \text{mm}^{-1}$

$T = 100$  K  
Block, colourless

$0.35 \times 0.29 \times 0.12$  mm

### Data collection

Bruker SMART APEXII DUO CCD area-  
detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(*SADABS*; Bruker, 2009)  
 $T_{\min} = 0.449$ ,  $T_{\max} = 0.735$

20560 measured reflections  
5389 independent reflections  
4508 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$   
 $\theta_{\max} = 32.5^\circ$ ,  $\theta_{\min} = 2.6^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -18 \rightarrow 18$   
 $l = -30 \rightarrow 28$

### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.064$   
 $S = 1.04$   
5389 reflections  
200 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0284P)^2 + 0.4844P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 0.50$  e  $\text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25$  e  $\text{\AA}^{-3}$

### Special details

**Experimental.** The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.04341 (2)	0.791941 (11)	0.448115 (6)	0.02747 (5)
F1	−0.59388 (13)	1.16479 (7)	0.96048 (4)	0.02911 (17)
O1	0.64441 (14)	0.86820 (8)	0.74360 (4)	0.02269 (17)
N1	0.18952 (16)	0.95514 (8)	0.82286 (5)	0.01709 (17)
N2	0.33651 (16)	0.94281 (8)	0.77529 (5)	0.01819 (18)
C1	−0.12723 (19)	1.02079 (10)	0.90775 (6)	0.0197 (2)
H1A	−0.0275	0.9644	0.9231	0.024*
C2	−0.2902 (2)	1.05229 (10)	0.94636 (6)	0.0219 (2)
H2A	−0.3024	1.0189	0.9883	0.026*
C3	−0.43476 (19)	1.13377 (10)	0.92222 (6)	0.0198 (2)
C4	−0.42365 (19)	1.18535 (9)	0.86209 (6)	0.0198 (2)
H4A	−0.5265	1.2404	0.8468	0.024*

C5	−0.25649 (19)	1.15407 (9)	0.82441 (6)	0.0182 (2)
H5A	−0.2431	1.1896	0.7832	0.022*
C6	−0.10798 (18)	1.07118 (9)	0.84631 (5)	0.01613 (19)
C7	0.06512 (18)	1.03787 (9)	0.80585 (5)	0.01641 (19)
C8	0.1174 (2)	1.09452 (10)	0.74299 (6)	0.0204 (2)
H8A	0.1713	1.1694	0.7527	0.024*
H8B	−0.0138	1.0972	0.7091	0.024*
C9	0.30095 (19)	1.02199 (9)	0.71935 (6)	0.0182 (2)
H9A	0.4379	1.0659	0.7171	0.022*
C10	0.23654 (18)	0.96573 (9)	0.65297 (5)	0.01642 (19)
C11	0.38005 (19)	0.96669 (9)	0.60390 (6)	0.0182 (2)
H11A	0.5182	1.0028	0.6123	0.022*
C12	0.3230 (2)	0.91522 (10)	0.54268 (6)	0.0193 (2)
H12A	0.4217	0.9153	0.5095	0.023*
C13	0.1196 (2)	0.86390 (9)	0.53106 (6)	0.0188 (2)
C14	−0.02684 (19)	0.86179 (10)	0.57892 (6)	0.0204 (2)
H14A	−0.1656	0.8263	0.5701	0.025*
C15	0.03330 (19)	0.91257 (10)	0.64006 (6)	0.0193 (2)
H15A	−0.0648	0.9111	0.6734	0.023*
C16	0.50858 (18)	0.87211 (10)	0.78447 (6)	0.0179 (2)
C17	0.5186 (2)	0.79944 (10)	0.84496 (6)	0.0226 (2)
H17A	0.6488	0.7524	0.8465	0.034*
H17B	0.3851	0.7546	0.8422	0.034*
H17C	0.5285	0.8441	0.8855	0.034*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.03694 (8)	0.02603 (7)	0.01781 (6)	−0.00295 (5)	−0.00372 (4)	−0.00310 (5)
F1	0.0272 (4)	0.0295 (4)	0.0332 (4)	0.0103 (3)	0.0145 (3)	0.0057 (3)
O1	0.0178 (4)	0.0292 (4)	0.0216 (4)	−0.0003 (3)	0.0047 (3)	−0.0060 (3)
N1	0.0168 (4)	0.0200 (4)	0.0149 (4)	0.0013 (3)	0.0036 (3)	−0.0005 (3)
N2	0.0192 (4)	0.0214 (4)	0.0148 (4)	0.0026 (4)	0.0052 (3)	0.0007 (3)
C1	0.0195 (5)	0.0191 (5)	0.0207 (5)	0.0037 (4)	0.0036 (4)	0.0029 (4)
C2	0.0220 (5)	0.0230 (5)	0.0217 (5)	0.0041 (4)	0.0068 (4)	0.0055 (4)
C3	0.0177 (5)	0.0195 (5)	0.0230 (5)	0.0014 (4)	0.0060 (4)	−0.0011 (4)
C4	0.0199 (5)	0.0161 (5)	0.0228 (5)	0.0031 (4)	0.0001 (4)	−0.0005 (4)
C5	0.0214 (5)	0.0162 (5)	0.0165 (5)	0.0011 (4)	0.0003 (4)	0.0004 (4)
C6	0.0167 (5)	0.0150 (4)	0.0165 (5)	−0.0004 (4)	0.0009 (4)	−0.0020 (4)
C7	0.0177 (5)	0.0164 (5)	0.0151 (5)	−0.0006 (4)	0.0019 (4)	−0.0011 (4)
C8	0.0266 (6)	0.0180 (5)	0.0171 (5)	0.0026 (4)	0.0051 (4)	0.0010 (4)
C9	0.0207 (5)	0.0184 (5)	0.0160 (5)	−0.0013 (4)	0.0038 (4)	0.0004 (4)
C10	0.0178 (5)	0.0162 (5)	0.0157 (5)	−0.0008 (4)	0.0037 (4)	0.0015 (4)
C11	0.0182 (5)	0.0195 (5)	0.0174 (5)	−0.0039 (4)	0.0044 (4)	0.0008 (4)
C12	0.0229 (5)	0.0202 (5)	0.0155 (5)	−0.0018 (4)	0.0055 (4)	0.0018 (4)
C13	0.0229 (5)	0.0174 (5)	0.0155 (5)	−0.0004 (4)	−0.0007 (4)	0.0006 (4)
C14	0.0172 (5)	0.0206 (5)	0.0231 (5)	−0.0025 (4)	0.0008 (4)	0.0008 (4)
C15	0.0173 (5)	0.0215 (5)	0.0199 (5)	−0.0017 (4)	0.0055 (4)	0.0010 (4)
C16	0.0157 (5)	0.0204 (5)	0.0173 (5)	−0.0007 (4)	0.0003 (4)	−0.0046 (4)
C17	0.0241 (5)	0.0243 (5)	0.0191 (5)	0.0057 (4)	0.0008 (4)	0.0001 (4)

*Geometric parameters (Å, °)*

Br1—C13	1.9003 (11)	C8—C9	1.5494 (16)
F1—C3	1.3622 (14)	C8—H8A	0.9900
O1—C16	1.2349 (14)	C8—H8B	0.9900
N1—C7	1.2919 (14)	C9—C10	1.5163 (16)
N1—N2	1.3947 (13)	C9—H9A	1.0000
N2—C16	1.3590 (15)	C10—C11	1.3945 (16)
N2—C9	1.4864 (15)	C10—C15	1.3987 (16)
C1—C2	1.3866 (16)	C11—C12	1.3940 (16)
C1—C6	1.4013 (16)	C11—H11A	0.9500
C1—H1A	0.9500	C12—C13	1.3867 (17)
C2—C3	1.3853 (16)	C12—H12A	0.9500
C2—H2A	0.9500	C13—C14	1.3887 (17)
C3—C4	1.3760 (17)	C14—C15	1.3916 (17)
C4—C5	1.3942 (17)	C14—H14A	0.9500
C4—H4A	0.9500	C15—H15A	0.9500
C5—C6	1.4008 (15)	C16—C17	1.5070 (17)
C5—H5A	0.9500	C17—H17A	0.9800
C6—C7	1.4653 (16)	C17—H17B	0.9800
C7—C8	1.5116 (16)	C17—H17C	0.9800
C7—N1—N2	107.94 (9)	N2—C9—C8	101.04 (9)
C16—N2—N1	121.60 (9)	C10—C9—C8	114.33 (10)
C16—N2—C9	124.40 (9)	N2—C9—H9A	109.8
N1—N2—C9	113.63 (9)	C10—C9—H9A	109.8
C2—C1—C6	120.72 (11)	C8—C9—H9A	109.8
C2—C1—H1A	119.6	C11—C10—C15	119.19 (10)
C6—C1—H1A	119.6	C11—C10—C9	120.12 (10)
C3—C2—C1	118.28 (11)	C15—C10—C9	120.69 (10)
C3—C2—H2A	120.9	C12—C11—C10	120.71 (11)
C1—C2—H2A	120.9	C12—C11—H11A	119.6
F1—C3—C4	118.69 (10)	C10—C11—H11A	119.6
F1—C3—C2	118.06 (11)	C13—C12—C11	118.83 (11)
C4—C3—C2	123.25 (11)	C13—C12—H12A	120.6
C3—C4—C5	117.78 (11)	C11—C12—H12A	120.6
C3—C4—H4A	121.1	C12—C13—C14	121.75 (11)
C5—C4—H4A	121.1	C12—C13—Br1	118.87 (9)
C4—C5—C6	121.09 (11)	C14—C13—Br1	119.37 (9)
C4—C5—H5A	119.5	C13—C14—C15	118.78 (11)
C6—C5—H5A	119.5	C13—C14—H14A	120.6
C5—C6—C1	118.87 (10)	C15—C14—H14A	120.6
C5—C6—C7	120.64 (10)	C14—C15—C10	120.73 (11)
C1—C6—C7	120.49 (10)	C14—C15—H15A	119.6
N1—C7—C6	120.77 (10)	C10—C15—H15A	119.6
N1—C7—C8	114.27 (10)	O1—C16—N2	120.09 (11)
C6—C7—C8	124.95 (10)	O1—C16—C17	123.28 (11)
C7—C8—C9	102.84 (9)	N2—C16—C17	116.61 (10)
C7—C8—H8A	111.2	C16—C17—H17A	109.5
C9—C8—H8A	111.2	C16—C17—H17B	109.5

C7—C8—H8B	111.2	H17A—C17—H17B	109.5
C9—C8—H8B	111.2	C16—C17—H17C	109.5
H8A—C8—H8B	109.1	H17A—C17—H17C	109.5
N2—C9—C10	111.65 (9)	H17B—C17—H17C	109.5
C7—N1—N2—C16	170.61 (10)	C16—N2—C9—C8	−168.23 (11)
C7—N1—N2—C9	−2.78 (13)	N1—N2—C9—C8	4.94 (12)
C6—C1—C2—C3	0.80 (19)	C7—C8—C9—N2	−4.85 (11)
C1—C2—C3—F1	−179.69 (11)	C7—C8—C9—C10	115.20 (10)
C1—C2—C3—C4	−0.44 (19)	N2—C9—C10—C11	−112.65 (12)
F1—C3—C4—C5	178.57 (10)	C8—C9—C10—C11	133.42 (11)
C2—C3—C4—C5	−0.67 (18)	N2—C9—C10—C15	67.46 (13)
C3—C4—C5—C6	1.44 (17)	C8—C9—C10—C15	−46.47 (15)
C4—C5—C6—C1	−1.10 (17)	C15—C10—C11—C12	−0.17 (17)
C4—C5—C6—C7	179.11 (11)	C9—C10—C11—C12	179.94 (10)
C2—C1—C6—C5	−0.05 (18)	C10—C11—C12—C13	0.69 (18)
C2—C1—C6—C7	179.74 (11)	C11—C12—C13—C14	−0.57 (18)
N2—N1—C7—C6	179.71 (10)	C11—C12—C13—Br1	−179.12 (9)
N2—N1—C7—C8	−0.93 (13)	C12—C13—C14—C15	−0.08 (18)
C5—C6—C7—N1	−173.72 (11)	Br1—C13—C14—C15	178.47 (9)
C1—C6—C7—N1	6.49 (17)	C13—C14—C15—C10	0.61 (18)
C5—C6—C7—C8	6.99 (17)	C11—C10—C15—C14	−0.49 (17)
C1—C6—C7—C8	−172.79 (11)	C9—C10—C15—C14	179.39 (11)
N1—C7—C8—C9	3.93 (13)	N1—N2—C16—O1	−175.15 (10)
C6—C7—C8—C9	−176.74 (10)	C9—N2—C16—O1	−2.50 (17)
C16—N2—C9—C10	69.83 (14)	N1—N2—C16—C17	6.25 (16)
N1—N2—C9—C10	−117.00 (10)	C9—N2—C16—C17	178.90 (10)

# Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C4—H4 <i>A</i> $\cdots$ O1 <sup>i</sup>	0.95	2.45	3.2772 (15)	146
C14—H14 <i>A</i> $\cdots$ F1 <sup>ii</sup>	0.95	2.50	3.3806 (15)	153
C15—H15 <i>A</i> $\cdots$ O1 <sup>iii</sup>	0.95	2.45	3.3800 (15)	166

Symmetry codes: (i)  $-x, y+1/2, -z+3/2$ ; (ii)  $-x-1, y-1/2, -z+3/2$ ; (iii)  $x-1, y, z$ .